



## Contributions of pollutants from North China Plain to surface ozone at the Shangdianzi GAW station

W. Lin, X. Xu, X. Zhang, J. Tang

### ► To cite this version:

W. Lin, X. Xu, X. Zhang, J. Tang. Contributions of pollutants from North China Plain to surface ozone at the Shangdianzi GAW station. *Atmospheric Chemistry and Physics Discussions*, 2008, 8 (3), pp.9139-9165. hal-00304169

**HAL Id: hal-00304169**

**<https://hal.science/hal-00304169>**

Submitted on 21 May 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.

# **Contributions of pollutants from North China Plain to surface ozone at the Shangdianzi GAW station**

**W. Lin<sup>1</sup>, X. Xu<sup>1</sup>, X. Zhang<sup>2</sup>, and J. Tang<sup>1</sup>**

<sup>1</sup>Key Lab. for Atmospheric Chemistry, Centre for Atmosphere Watch and Services, Chinese Academy of Meteorological Sciences, China Meteorological Administration, Beijing, China

<sup>2</sup>Institute of Urban Meteorology, China Meteorological Administration, Beijing, China

Received: 29 February 2008 – Accepted: 1 April 2008 – Published: 21 May 2008

Correspondence to: W. Lin (linwl@cams.cma.gov.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Regional ozone pollution has become one of the top environmental concerns in China, especially in those economically vibrant and densely populated regions, such as North China region including Beijing. To address this issue, surface ozone and ancillary data over the period 2004–2006 from the Shangdianzi Regional Background Station in north China were analyzed. Due to the suitable location and valley topography of the site, transport of pollutants from the North China Plain was easily observed and quantified according to surface wind directions. Regional (polluted) and natural (clean) background ozone concentrations were obtained by detailed statistic analysis. Contribution of pollutants from North China Plain to surface ozone at SDZ was estimated by comparing ozone concentrations observed under SW wind conditions and that under NE wind conditions. The average daily accumulated ozone contribution was estimated to be 240 ppb-hr. The average regional contributions to surface ozone at SDZ from the North China Plain were 21.8 ppb for the whole year, and 19.2, 28.9, 25.0, and 10.0 ppb for spring, summer, autumn, and winter, respectively. The strong ozone contribution in summer led to disappearance of the spring ozone maximum phenomenon at SDZ under winds other than from the WNN to E sectors. High winter NO<sub>x</sub> concentrations in the North China Plain caused negative ozone contribution in winter.

## 1 Introduction

Being one of the greenhouse gases and a key oxidant, ozone plays important role in atmospheric energy budget and chemistry. Changes in atmospheric ozone may contribute to climate change and high levels of surface ozone can cause adverse effect on the health of human being and ecosystem. Anthropogenic emissions have led to significant changes in atmospheric ozone. These changes, including the increase in tropospheric ozone (Guicherit and Roemer, 2000; Vingarzan, 2004) and the depletion of stratospheric ozone (UNEP, 1989, 1994, 1998) have been important factors for cli-

ACPD

8, 9139–9165, 2008

### Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

mate change (Fortster et al., 2007). In the troposphere, photolysis of ozone by solar UV radiation to electronically excited  $O(^1D)$  and the subsequent reaction with water vapor is the major source of OH radical. OH is one of the key species for the chemical reactions in the atmosphere and its abundance is an important index of oxidizing capacity of the atmosphere (Kley et al., 1999; Thompson, 1992; Jacob, 2003). At high pH value ( $pH > 5$ ), ozone in aqueous reacts rapidly with  $S(IV)$  to form sulfate and hence contributes to acid deposition (Tanner and Schorran, 1995).

Ozone originates from in-situ photochemical production in the reactions of its precursors ( $NO_x$ , CO, VOCs, and etc.) and from vertical and horizontal transport. On the scale of 100 years, the increasing trends of tropospheric ozone are qualitatively in agreement with emission trends of precursors and the increase in tropospheric ozone level has become one of the most crucial environmental problems to solve in the coming decades (Finlayson-Pitts and Pitts, 1997; Guicherit and Roemer, 2000). Ozone at ground level is usually much more influenced by emissions. Because natural and anthropogenic ozone precursor emissions vary strongly from region to region (Staehlin et al., 2001), photochemical sources of ozone have high regional variability. In many mega cities and polluted areas (e.g., city clusters in China and other countries), serious ozone pollution caused by large precursor emissions has concerned civilians and decision-makers (Molina and Molina, 2004; Shao et al., 2006).

China has experienced unprecedented economic growth over the past two decades, accompanied by the development of large-scale industries and services. However, the economic boom has led to a general decline in environmental quality, especially in urban city and in city cluster regions (Zhang et al., 1998; Hao and Wang, 2005; Shao et al., 2006). More and more concerns have been focused on the regional pollutions in those leading economic regions, for example, the Yangtze River delta (Wang et al., 2005), the Pearl River delta (Wang et al., 2003), and the Beijing–Tianjin–Bohai Bay (Xu et al., 2005; Ding et al., 2008). In China, most measurements of surface ozone have been done in urban areas. Since a few years, instruments for surface ozone and other pollutants have been set up at China's background stations to monitor the long-

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

term changes in these pollutants over a regional scale. Some data obtained have been used to study the long-term trends of surface ozone at a regional background station in eastern China (Xu et al., 2008). Ozone and related gases have been measured continuously since 2004 at Shangdianzi (SDZ) regional background station. In this paper, we present analysis of surface ozone data from this station and focus on the contributions of pollutants from North China Plain to natural background level of surface ozone at the site. A more general analysis of the reactive gases data from the station is given in another paper<sup>1</sup>.

## 2 Site and observations

Shangdianzi (SDZ; 40°39'N, 117°07'E, 293.9 m a.s.l.) station is one of the regional Global Atmosphere Watch (GAW) stations in China. The station is located in the northern part of North China Plain and in the Miyun County of Beijing. It is about 100 km and 55 km northeasterly to the urban area and Miyun Township of Beijing, respectively (see Fig. 1). Within 30 km of the site, there are only small villages with sparse population and thus insignificant anthropogenic emission sources. The instrument building of the station is situated on the south slope of a hill, which is surrounded by mountainous areas except the southwest direction. Due to the valley topography, the prevailing winds are northeasterly and southwesterly as shown in Fig. 2. Therefore, polluted air masses from urban areas and satellite towns of Beijing can be easily transported to SDZ under southwesterly winds, while relatively clean air masses under other winds.

A set of commercial instruments from Thermo Environmental Instruments, Inc., USA has been used to measure O<sub>3</sub> (TE 49C), NO/NO<sub>2</sub>/NO<sub>x</sub> (TE 42CTL), CO (TE 48C), and SO<sub>2</sub> (TE 43CTL) since January, 2004. Daily zero/span checks are automatically done using a dynamic gas calibrator (TE 146C) in combination with a zero air supply (TE

<sup>1</sup> Meng, Z., Xu, X., Lin, W., Ding, G., Tang, J., Yan, P.: Characteristics of trace gaseous pollutants at a regional background station in Northern China, submitted to Atmos. Chem. Phys., 2008.

### Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



111) and a set of standard reference gas mixtures (Chemical Metrology and Analytical Science Division, National Institute of Metrology, Beijing, China). All instruments are housed in an air-conditioned room. Measurement signals are acquired every 6 s and recorded as 5-min averages. Multipoint calibrations are made every 3–6 months.

5 For ozone, the standards are traceable to the Standard Reference Photometer (SRP) maintained by WMO World Calibration Centre in Switzerland (EMPA). The national standard gases of SO<sub>2</sub>, NO, and CO are compared against NIST-traceable standards from Scott Specialty Gases, USA. After the correction of data on the basis of multipoint calibration, hourly average O<sub>3</sub> data are calculated and are used for further analysis  
10 in this paper. Meteorological data, including wind, temperature, relative humidity, and etc., are also obtained from the site, with a resolution of 1 h.

### 3 Results and discussion

#### 3.1 Diurnal and seasonal variations

15 Figure 3 shows the average diurnal variations in each month and monthly variations of surface O<sub>3</sub> and O<sub>x</sub> (O<sub>3</sub>+NO<sub>2</sub>) concentrations at SDZ. Some detailed statistics of O<sub>3</sub> and O<sub>x</sub>, including the monthly-averaged diurnal maxima/minima with their appearing time, diurnal amplitudes, and the monthly mean concentrations in each month are listed in Table 1. As can be seen in Fig. 3, both O<sub>3</sub> and O<sub>x</sub> concentrations show typical  
20 diurnal cycles related to photochemistry, with maxima in the afternoon and minima in the early morning. The ozone maxima were in late afternoon (15:00–18:00) and the minima were in the morning (05:00–07:00). The higher value the ozone maxima kept, the later the appearing time is. The diurnal variations of both quantities in winter are much less significant than those in other seasons. The highest diurnal amplitude of ozone appears in June. So does the highest monthly mean ozone concentration. In  
25 September, in which the monthly mean ozone concentration shows a secondary peak, the diurnal amplitude of ozone is also high and only next to the highest one.

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The diurnal amplitudes of surface ozone at SDZ are much more pronounced than those at the Waliguan (36.30° N, 100.90° E, 3810 m a.s.l.) Global Baseline Station, a Chinese GAW station at a remote mountain site, where the ozone level is closer to natural background, with small diurnal variations (Ma et al., 2002a; Wang et al., 2006).

5 The diurnal pattern and variability of surface ozone depend very much on the intensity of UV radiation, level and variations of ozone precursors, and vertical and horizontal transport of ozone. In situ photochemical production does not significantly contribute to the diurnal variation of surface ozone at Waliguan because of the extremely low levels of ozone precursors, while at Linan and Longfengshan, the regional GAW stations in  
10 eastern and northeastern China, photochemistry plays a key role in diurnal variation of surface ozone (Ma et al., 2002b). Compared to Waliguan, SDZ is much closer to large cities like Beijing and hence larger emission sources of ozone precursors. Photochemistry at this site should be much more intensive. This can explain the difference in the diurnal variability of surface ozone between the two stations.

15 As shown in Table 1, the monthly mean ozone concentration is much lower in the winter months than in the other months. For example, the monthly mean ozone concentration in December is only 17.7 ppb compared to 46.5 ppb in June. Although the monthly mean  $O_x$  concentration is also low in winter months, the summer-winter difference in the  $O_x$  concentration is by far not as large as that in the ozone concentration. In  
20 other words, from summer to winter, the total oxidant, as estimated by  $O_3 + NO_2$ , drops less than the ozone concentration. This is because  $NO_2$  contributed about 48% to  $O_x$  in winter (52% in December) and only 10% in June.

25 The year round and summertime average diurnal variations of ozone and related gases are shown in Fig. 4. The ozone concentration increases from its minimum in early morning to its maximum in late afternoon and then decreases till next morning. In recent years, ozone in Beijing city peaks around noon because of photochemical production under the daily strong solar radiation and abundant precursors (Shao et al., 2006). However, as shown in Fig. 4, the ozone concentration in SDZ peaks near dusk and continues its increase in the afternoon, accompanied with the increase of  $NO_x$  and

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

CO concentrations. This indicates transport of ozone and other pollutants from urban area to SDZ. The strong influence of air quality in rural area by urban plume of Beijing has been also observed by Wang et al. (2006) and Zheng et al. (2005).

### 3.2 Ozone frequency distribution

5 Surface ozone frequency distribution (4 ppb per bin) with Lorentz distribution curve fitting is shown in Fig. 5. The total number of hourly average data is 25356, with the mean  $\pm$ SD of 31.5 $\pm$ 19.9 ppb, median of 29.4 ppb, 25% percentile of 17.7 ppb, 75% percentile of 41.0 ppb, maximum of 155.7 ppb, and minimum of 0 ppb. The peak concentration in Fig. 5 is equal to the mean value and close to the median value. Since  
10 the concentration of surface ozone at any site and any time depends on the dynamic balance between ozone producing and destroying processes and transport conditions. To some extent, the peak concentration of the ozone frequency distribution, which appears at the highest probability, represents well the local background level of surface ozone. Therefore, average value in each month could stand for the monthly ozone  
15 background level.

### 3.3 Dependence of surface ozone level on wind direction

Since wind direction gives an insight into the first order advection processes, its correlation with observed ozone data can provide the information about the impact of air-mass transport on the surface ozone variability. In this analysis, the wind direction is divided  
20 into 16 standard sectors (clockwise from North) and ozone and  $O_x$  concentrations in the corresponding sectors are averaged. Figure 6 shows the rose distributions of the average  $O_3$  and  $O_x$  concentrations and  $O_3$  concentration under calm wind conditions (wind speed  $<0.3$  m/s). The rose distributions of ozone and  $O_x$  indicate that average ozone and  $O_x$  concentrations were higher in the SW-SWW-W sectors than in other  
25 sectors. The lowest ozone and  $O_x$  levels were in the NEE sector. The average ozone concentration in the NEE sector was close to that under calm wind conditions (with a

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



difference of less than 2 ppb) and 21.8 ppb lower than that in the SWW sector.

Figure 7 shows the average  $O_3$  distribution in different wind direction sectors in four seasons. Except in winter, the average  $O_3$  concentrations in the SW-SWW-W sectors were significantly higher than in other sectors. The lowest concentration was in the NEE sector. The highest  $O_3$  concentration in winter appeared in the N sector. The spring ozone maximum is a Northern Hemispheric phenomenon (Monks, 2000; Vingarzan, 2004). At SDZ, however, the springtime ozone maximum appeared only in the WNN to E sectors. For the other sectors, highest ozone concentrations appeared in summer and the ozone levels in spring were a little lower than those in summer. We believe that air masses transported over the polluted North China Plain had produced more ozone in summer than in any other seasons, leading to the disappearance of the spring ozone maximum in the other sectors. Table 2 summarizes the maximum ([Max]) and minimum ([Min]) values of the annual and seasonal average ozone concentrations and the corresponding appearing sectors. Average ozone concentrations under calm wind conditions ([CW]) and the differences between [Max] and [Min] and between [Max] and [CW] are also shown in the table. The ozone difference between [Max] and [Min] was 21.8 ppb for the whole year, and was 19.2, 28.9, 25.0, and 10.0 ppb in spring, summer, autumn, and in winter, respectively.

Figure 8 shows the rose distributions of seasonal average  $SO_2$ ,  $NO_x$ , CO, temperature, relative humidity (RH), and wind speed. As can be seen in Fig. 8a, b, and c, in all seasons except summer, the levels of  $SO_2$ ,  $NO_x$ , and CO concentrations in the S-W sectors are much higher than in other sectors, while those in the N-NEE sectors much lower than in other sectors, indicating that polluted air masses are mainly from the S-W sectors (i.e., the North China Plain) and clean air masses are mainly from the N-NEE sectors. The rose distribution of RH shown in Fig. 8d indicates that except in summer, air masses from the N sector are significantly drier than from other sectors. Because air masses from the N sector are characterized as having low humidity, relatively high ozone, and lower  $SO_2$ ,  $NO_x$ , and CO concentrations, it is likely that they contain more air from the free troposphere than those from other sectors. The rose distributions

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of temperature and wind speed, as shown in Fig. 8e and f, respectively, suggest that air masses from the North China Plain are warmer and are transported to SDZ under higher local wind speed conditions.

Observed ozone concentrations in air from different sectors (clean and polluted) should stand for the levels of different background ozone. To simplify the situation concerned here, we differentiate two kinds of background ozone, natural background ozone and regional background ozone, following the definitions by Kourtidis et al. (1997) and Bronnimann et al. (1998). Natural background ozone means ozone generated chemically in the troposphere from biogenic or geogenic emissions plus ozone transported from the stratosphere. Regional background ozone is the ozone level over a large area ( $1000 \times 1000 \text{ km}^2$ ) produced by the mixing of air masses of different origin outside and inside the defined area. The definition of regional background is made under the ideal status of “well mixing”. So far, it is difficult to recognize the exact regional background ozone for a certain moment. In view of the fact that SDZ is quite distant from major sources, we assume that air masses from the North China Plain have been well mixed before arriving at SDZ. Under this assumption, the average of the surface ozone concentration in the southwest sector for a given period can be considered as the typical value of the regional (polluted) background ozone of the North China Plain for this period. Ozone in air masses from the clean sectors, i.e., the N-NEE sectors can be treated as the natural background ozone for SDZ. Therefore, the contribution of pollutants from the North China Plain to the ozone level at SDZ can be estimated to be the difference between the regional background and the natural background ozone for the site.

Figure 9 shows the year round average diurnal variations of ozone concentrations corresponding to the N-E and S-W wind direction sectors. The difference between the S-W and N-E curves is also shown as bars in Fig. 9. Although the two diurnal variation curves look similar in terms of their general patterns, they differ at points. The first and profound difference is that the S-W curve shows higher ozone concentrations than the N-E curve at all time of the day, particularly in the afternoon. The second difference

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

is the timing of the maximum and minimum ozone concentrations, with the maximum and minimum in the S-W curve being 1–2 h later than those in the N-E curve. The S-W curve has the typical features of the diurnal ozone variation in aged air masses, e.g., enhanced ozone level and later ozone daily maximum. This supports the idea that ozone concentration at SDZ under the southwesterly wind condition is very strongly influenced by the transport of polluted air masses from the North China Plain and photochemical production during the transport of the air masses. Assuming that the S-W and N-E curves represent the average diurnal variations of the regional background ozone and the natural background ozone at SDZ, respectively, we can estimate the net contribution of the North China Plain to surface ozone at SDZ using the difference between the S-W and N-E curves. As shown in Fig. 9, the difference between the two curves is at highest 20 ppb (17:00) and at lowest 1.6 ppb (07:00). The daily average of the difference is 10 ppb, which gives an accumulated daily contribution of 240 ppb-hr.

The ozone contribution of the North China Plain depends highly on season. Figure 10 shows the seasonal variations of ozone concentrations corresponding to the S-W and N-E wind direction sectors and their difference. Similar to diurnal variation in Fig. 9, the seasonal variations of the ozone difference between the S-W and N-E curves in Fig. 10 can be used to estimate the net contribution of the North China Plain to surface ozone at SDZ for the time scale of month, season, and year. There are two peaks in the seasonal variation of the ozone difference, with the primary one (about 34 ppb) being in September and the secondary one (about 29 ppb) being in June. In winter, the ozone difference is negative, suggesting that air masses from the North China Plain contain less ozone than those from the N-E sector. This does not mean that the winter air in the North China Plain is cleaner than the natural background at all. Rather it is caused by the pollution in the North China Plain. Because of the inactivity of photochemical production in winter, much higher  $\text{NO}_x$  concentrations in the SW sectors (see Fig. 8c) may consume (through “gas titration”) a portion of ozone in the natural background, leading to the negative values shown in Fig. 10. To prove this, the seasonal variations of  $\text{O}_3$  corresponding to the S-W and N-E wind direction sectors

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and their difference are plotted in Fig. 11. Obviously, the  $O_x$  difference is positive in all seasons, suggesting that even in the cold season the total oxidant is not reduced by the pollutants from the North China Plain, but enhanced.

## 4 Conclusions

5 Measurements of surface ozone at SDZ from the period 2004–2006, together with ancillary data, were summarized and used to study the impact of pollutants from the North China Plain on surface ozone at the background site. Influenced by the location and topography of the SDZ site, surface ozone there has something unique. The seasonal variation of surface ozone at SDZ shows a double-peak pattern, with the primary and  
10 secondary peaks being in June and September, respectively. This pattern reflects the effect of Asian summer monsoon on the level of surface ozone at the site. Monthly mean concentrations of surface ozone from January to December were 20.3, 27.0, 35.0, 40.1, 41.4, 46.5, 33.6, 30.9, 34.7, 28.3, 19.4, and 17.7 ppb, respectively. The average diurnal variation of surface ozone at SDZ takes the form of single wave and  
15 shows significant influence of photochemistry. An unusual phenomenon is that ozone peaks near dusk and continues its increase in the afternoon, accompanied with the increase of  $NO_x$  and CO concentrations. This can be attributed to transport of ozone and other pollutants from urban area to SDZ.

The frequency distribution of hourly mean ozone concentration at SDZ shows that  
20 the most probable ozone concentration is equal to the mean value and close to the median value. Therefore, the average value in each month may stand for the monthly ozone background level. Being a station at the north edge of the North China Plain and a site with valley topography, SDZ is an ideal site to be used to capture the regional (polluted) background of North China Plain and the natural (clean) background according to different wind directions. Contribution of pollutants from North China Plain to  
25 surface ozone at SDZ can be estimated by comparing ozone concentrations observed under SW wind conditions and that under NE wind conditions. On the average, the

### Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

daily accumulated ozone contribution amounts to 240 ppb-hr. The average regional contributions to surface ozone at SDZ from the North China Plain are estimated to be 21.8 ppb for the whole year, and 19.2, 28.9, 25.0, and 10.0 ppb for spring, summer, autumn, and winter, respectively. From May to September, the estimated ozone contributions are all above 20 ppb with the most significant contribution (34 ppb) being in September. Caused by the strong ozone contribution in summer, the spring ozone maximum phenomenon disappears at SDZ under winds other than from the WNN to E sectors. The ozone contribution in winter is negative due to impact of high NO<sub>x</sub> concentrations in the North China Plain.

*Acknowledgements.* We thank the staff of the Shangdianzi station for carrying out the measurements. This work is supported by the “National Basic Research Program of China” (No. 2005CB4222002) and by Chinese Natural Science Foundation (Grants 20407001 and 40775074) and by supported by the CMA project (CCSF2007-15). X. Xu thanks the grant from the Ministry of Personnel of the Peoples Republic of China.

## References

- Bronnimann, S., Schuepbach, E., Zanis, P., Buchmann, B., and Wanner, H.: A climatology of regional background ozone at different elevations in Switzerland (1992–1998), *Atmos. Environ.*, 34, 5191–5198, 2000.
- Ding, A. J., Wang, T., Thouret, V., Cammas, J.-P., and Nedelec, P.: Tropospheric ozone climatology over Beijing: analysis of aircraft data from the MOZAIC program, *Atmos. Chem. Phys.*, 8, 1–13, 2008, <http://www.atmos-chem-phys.net/8/1/2008/>.
- Finlayson-Pitts, B. J. and Pitts, J. J. N.: Tropospheric air pollution: ozone, airborne toxics, polycyclic aromatic hydrocarbons, and particles, *Science*, 276, 1045–1051, 1997.
- Forster P., Ramaswamy V., Artaxo P., et al.: Changes in Atmospheric Constituents and in Radiative Forcing. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B.,

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- Guicherit, R. and Roemer, M.: Tropospheric ozone trends, *Chemosphere: Global Change Science*, 2, 167–183, 2000.
- 5 Hao, J. M. and Wang, L.T.: Improving urban air quality in China: Beijing case study, *J. Air Waste Manage. Assoc.*, 55, 1298–1305, 2005.
- Jacob, D. J.: The Oxidizing Power of the Atmosphere, in: Potter T, Colman B, Fishman J, et al. *Handbook of Weather, Climate and Water*, Chichester: John Wiley & Sons, Inc., 1–18, 2003.
- Kley, D., Kleinmann, M., Sanderman, H., Krupa, S.: Photochemical oxidants: state of the science, *Environmental Pollution*, 100, 19–42, 1999.
- 10 Kourtidis, K., Ziomas, I., Zerefos, C., Balis, D., Suppan, P., Vasaras, A., Kosmidis, V., and Kentarchos, A.: On the background ozone values in Greece, in: *The Oxidizing Capacity of the Troposphere*, edited by: Larsen, B., Versino, B., Angeletti, G., EUR 17482, EU, Brussels, 387–390, 1997.
- 15 Ma, J., Tang, J., Zhou, X., and Zhang, X.: Estimates of the chemical budget for ozone at Waliguan Observatory, *J. Atmos. Chem.*, 41, 21–48, 2002a.
- Ma, J., Zhou, X., and Hauglustaine, D.: Summertime tropospheric ozone over China simulated with a regional chemical transport model: 2. Source contributions and budget, *J. Geophys. Res.*, 107(D22), 4612, doi:10.1029/2001JD001355, 2002b.
- 20 Molina, M. J. and Molina, L. T.: Megacities and atmospheric pollution, *J. Air Waste Manage. Assoc.*, 54, 644–680, 2004.
- Monks, P. S.: A review of the observations and origins of the spring ozone maximum, *Atmos. Environ.*, 34, 3545–3561, 2000.
- Shao, M., Tang, X., Zhang, Y., and Li, W.: City clusters in China: air and surface water pollution, *Front Ecol Environ*, 4(7), 353–361, 2006.
- 25 Staehlin, J., Harris, N. R. P., Appenzeller, C., and Eberhard, J.: Ozone Trends: a Review, *Reviews of Geophysics*, 39(2), 231–290, 2001.
- Tanner, R. L. and Schorran, D. E.: Measurement of gaseous peroxides near the Grand Canyon, implication for summertime visibility impairment from aqueous phase secondary sulfate formation, *Atmos. Environ.*, 29, 1113–1122, 1995.
- 30 Thompson, A.: *The Oxidizing Capacity of the Earth's Atmosphere: Probable Past and Future Changes*, Science, 256, 1157–1165, 1992.
- UNEP.: *Environmental Effects Panel Report*, Nairobi: UNEP, 1989.

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



UNEP: Environmental Effects of Ozone Depletion: 1994 and 1998 Assessment, Nairobi: UNEP, 1994 and 1998.

Vingarzan, R.: A review of surface ozone background levels and trends, *Atmos. Environ.*, 38: 3431–3442, 2004.

5 Wang, H., Kiang, C.S., Tang, X., Zhou, X., and Chameides, W.: Surface ozone: A likely threat to crops in Yangtze delta of China, *Atmos. Environ.*, 39, 3843–3850, 2005.

Wang, T., Ding, A., Gao, J., and Wu, W.: Strong ozone production in urban plumes from Beijing, China, *Geophys. Res. Lett.*, 33, L21806, doi:10.1029/2006GL027689, 2006.

10 Wang, T., Poon, C. N., Kwok, Y. H., and Li, Y. S.: Characterizing the temporal variability and emission patterns of pollution plumes in the Pearl River Delta of China, *Atmos. Environ.*, 37, 3539–3550, 2003.

Wang, T., Wong, H., Tang, J., Ding, A., Wu, W. and Zhang, X.: On the origin of surface ozone and reactive nitrogen observed at a remote mountain site in the north-eastern Qinghai-Tibetan Plateau, western China, *J. Geophys. Res.*, 111, D08303, doi:10.1029/2005JD006527, 2006.

15 Xu, X., Lin, W., Wang, T., Yan, P., Tang, J. Meng, Z., and Wang, Y.: Long-term trend of surface ozone at a regional background station in eastern China 1991–2006: Enhanced variability, *Atmos. Chem. Phys. Discuss.*, 8, 215–243, 2008.

20 Xu, X., Xie, L., Ding, G., and Bian, L.: Beijing Air Pollution Project To Benefit 2008 Olympic Games, American Meteorological Society, 1543–1544, 2005.

Zhang, Y., Shao, K., Tang, X., and Li, J.: The Study of Urban Photochemical Smog Pollution in China, *Acta Scientiarum Naturalium Universitatis Pekinensis* (Chinese with English abstract), 34(2–3), 392–400, 1998.

25 Zheng, X., Ding, G., Yu, H., Liu, Y., and Xu, X.: Vertical distribution of ozone in the planetary boundary layer at the Ming Tombs, Beijing, *Science in China (D Earth Sciences)*, 48 (Supp.), 55–63, 2005.

ACPD

8, 9139–9165, 2008

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** The average diurnal maxima/minima with their appeared time, the diurnal ranges, and monthly means of ozone at SDZ.

Month	Mean	O <sub>3</sub> (ppb)			Mean	O <sub>x</sub> (ppb)		
		Diurnal Max (appearing time)	Diurnal Min (appearing time)	Diurnal amplitude		Diurnal Max (appearing time)	Diurnal Min (appearing time)	Diurnal amplitude
Jan	20.3	29.4 (16:00)	14.6(07:00)	14.8	35.9	40.9(17:00)	30.0(09:00)	10.9
Feb	27.0	35.7(15:00)	20.5(02:00)	15.2	37.6	44.5(17:00)	31.3(08:00)	13.1
Mar	35.0	45.8(17:00)	23.3(07:00)	22.5	44.1	53.6(17:00)	34.7(07:00)	18.9
Apr	40.1	54.8(16:00)	24.3(06:00)	30.5	48.7	62.4(16:00)	35.5(07:00)	26.8
May	41.4	58.6(18:00)	23.9(06:00)	34.7	49.1	65.5(18:00)	34.1(06:00)	31.4
Jun	46.5	69.2(18:00)	23.7(06:00)	45.5	51.7	73.5(18:00)	31.2(06:00)	42.3
Jul	33.6	50.3(16:00)	15.0(06:00)	35.4	39.2	54.6(18:00)	22.1(06:00)	32.5
Aug	30.9	49.5(17:00)	13.1(06:00)	36.4	37.7	54.5(17:00)	21.5(06:00)	33.0
Sep	34.7	57.4(17:00)	15.7(06:00)	41.7	47.8	67.2(17:00)	31.0(07:00)	36.2
Oct	28.3	46.6(16:00)	13.5(06:00)	33.1	46.2	61.5(17:00)	32.4(07:00)	29.1
Nov	19.4	31.9(15:00)	11.0(06:00)	20.9	36.5	43.4(17:00)	27.6(08:00)	15.8
Dec	17.7	24.9(15:00)	13.5(06:00)	11.4	36.9	41.9(17:00)	30.1(09:00)	11.8

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.

**Table 2.** The annual and seasonal statistics of ozone concentrations and the corresponding appearing sectors.

Period	[Max]*		[Min]*		[CW]*	[Max]-[Min]	[Max]-[CW]
Year	43.7	SW	21.9	NEE	20.2	21.8	23.5
MAM	48.4	SW	28.2	NEE	22.7	20.2	25.8
JJA	52.1	SWW	23.2	NEE	22.9	28.9	29.2
SON	43.0	SWW	18.0	NEE	20.8	25.0	22.2
DJF	28.3	N	18.3	W	12.4	10.0	15.9

\*: [Max]=maximum average;


[Min]=minimum average;

[CW]=average under calm wind conditions

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Contributions to surface ozone at Shangdianzi GAW station

W. Lin et al.



**Fig. 1.** Map showing the location of the SDZ station (white cross), topography and land use in the surrounding area. Map from <http://worldwind.arc.nasa.gov>.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion


**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.

**Fig. 2.** Wind direction rose of SDZ (from year 2004 to 2006).[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.




**Fig. 3.** Average diurnal variations in each month and variations of monthly mean surface  $O_3$  and  $O_x$  concentrations at SDZ.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.



**Fig. 4.** Average diurnal variations of  $O_3$ ,  $NO/NO_2/NO_x$ ,  $SO_2$ , and  $CO$  at SDZ: the whole year (up), summer (bottom).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.




**Fig. 5.** Surface ozone frequency distribution (4 ppb for each bin) with Lorentz distribution curve fitting at SDZ.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.




**Fig. 6.** Average  $O_3$  and  $O_x$  distributions according to wind directions and average  $O_3$  concentration under calm wind conditions ( $O_3$ -CW). Unit: ppb.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**


W. Lin et al.

**Fig. 7.** Seasonal average ozone concentrations in different wind direction sectors.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.



**Fig. 8.** Dependence of seasonal average  $\text{SO}_2$ ,  $\text{NO}_x$ , CO, temperature, relative humidity (RH), and wind speed (WS) on wind direction.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.




**Fig. 9.** The annually averaged diurnal variations of ozone concentrations for N-E and S-W wind direction sectors.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.




**Fig. 10.** Seasonal variations of average  $O_3$  concentrations corresponding to the S-W and N-E wind direction sectors and their difference.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contributions to  
surface ozone at  
Shangdianzi GAW  
station**

W. Lin et al.



**Fig. 11.** Seasonal change of average  $O_x$  concentrations corresponding to the S-W and N-E wind direction sectors and their difference.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)